

Quasinormal modes of a black hole in the deformed Hořava-Lifshitz gravity

Songbai Chen^{*} and Jiliang Jing[†]

*Institute of Physics and Department of Physics,
Hunan Normal University, Changsha, Hunan 410081, P. R. China
Key Laboratory of Low Dimensional Quantum Structures and Quantum
Control (Hunan Normal University), Ministry of Education, P. R. China.*

Abstract

We study the quasinormal modes of the massless scalar perturbation in the background of a deformed black hole in the Hořava-Lifshitz gravity with coupling constant $\lambda = 1$. Our results show that the quasinormal frequencies depend on the parameter in the Hořava-Lifshitz gravity and the behavior of the quasinormal modes is different from those in the Reissner-Norström and Einstein-Born-Infeld black hole spacetimes. The absolute value of imaginary parts is smaller and the scalar perturbations decay more slowly in the deformed Hořava-Lifshitz black hole spacetime. These information can help us understand more about the Hořava-Lifshitz gravity.

PACS numbers: 04.30.-w, 04.62.+v, 97.60.Lf.

^{*}Electronic address: csb3752@163.com

[†]Electronic address: jljing@hunnu.edu.cn

Inspiring by the Lifshitz model, Hořava [1] proposes recently a field theory model for a UV complete theory of gravity, which is a non-Lorentz invariant theory of gravity in $3 + 1$ dimensions. Unlike Einstein gravity, it is renormalizable by power-counting arguments. Thus, it is believed widely that it could be a candidate for Einstein's general relativity. Very recently, the Hořava-Lifshitz gravity theory has been intensively investigated in [2–11] and its cosmological applications have been studied in [12–18]. Some static spherically symmetric black hole solutions have been found in Hořava-Lifshitz theory [19–24] and the associated thermodynamic properties with those black hole solutions have been investigated in [25–28].

The four-dimensional metric in the ADM formalism can be expressed as [29]

$$ds_{ADM}^2 = -N^2 dt^2 + g_{ij} \left(dx^i - N^i dt \right) \left(dx^j - N^j dt \right) \quad (1)$$

and the Einstein-Hilbert action is given by

$$S_{EH} = \frac{1}{16\pi G} \int d^4x \sqrt{g} N \left(K_{ij} K^{ij} - K^2 + R - 2\Lambda \right), \quad (2)$$

where G is Newton's constant and K_{ij} is extrinsic curvature which takes the form

$$K_{ij} = \frac{1}{2N} \left(\partial_t g_{ij} - \nabla_i N_j - \nabla_j N_i \right). \quad (3)$$

In general, the IR vacuum of this theory is anti de Sitter (AdS) spacetimes. To obtain a Minkowski vacuum in the IR sector, one can modify the theory by introducing “ $\mu^4 R$ ” and then take the $\Lambda_W \rightarrow 0$ limit. This does not change the UV properties of the theory, but it alters the IR properties. The deformed action of the non-relativistic renormalizable gravitational theory is given by [20]

$$S_{HL} = \int dt d^3x, \left(\mathcal{L}_0 + \tilde{\mathcal{L}}_1 \right) \quad (4)$$

$$\mathcal{L}_0 = \sqrt{g} N \left\{ \frac{2}{\kappa^2} (K_{ij} K^{ij} - \lambda K^2) + \frac{\kappa^2 \mu^2 (\Lambda_W R - 3\Lambda_W^2)}{8(1-3\lambda)} \right\}, \quad (5)$$

$$\tilde{\mathcal{L}}_1 = \sqrt{g} N \left\{ \frac{\kappa^2 \mu^2 (1-4\lambda)}{32(1-3\lambda)} R^2 - \frac{\kappa^2}{2w^4} \left(C_{ij} - \frac{\mu w^2}{2} R_{ij} \right) \left(C^{ij} - \frac{\mu w^2}{2} R^{ij} \right) + \mu^4 R \right\}. \quad (6)$$

Here C_{ij} is the Cotton tensor, defined by

$$C^{ij} = \epsilon^{ik\ell} \nabla_k \left(R^j_\ell - \frac{1}{4} R \delta^j_\ell \right) = \epsilon^{ik\ell} \nabla_k R^j_\ell - \frac{1}{4} \epsilon^{ikj} \partial_k R. \quad (7)$$

Comparing the action to that of general relativity in the ADM formalism, one can find that the speed of light, Newton's constant and the cosmological constant are given by

$$c = \frac{\kappa^2 \mu}{4} \sqrt{\frac{\Lambda_W}{1-3\lambda}}, \quad G = \frac{\kappa^2}{32\pi c}, \quad \Lambda = \frac{3}{2} \Lambda_W. \quad (8)$$

Taking $N^i = 0$, the spherically symmetric solutions could be obtained with the metric ansatz [19, 21–27]

$$ds^2 = -N^2(r) dt^2 + \frac{dr^2}{f(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (9)$$

Substituting the metric ansatz (9) into the action, and then vary the functions N and f , one can find that the reduced Lagrangian reads

$$\tilde{\mathcal{L}}_1 = \frac{\kappa^2 \mu^2 N}{8(1-3\lambda)\sqrt{f}} \left(\frac{\lambda-1}{2} f'^2 - \frac{2\lambda(f-1)}{r} f' + \frac{(2\lambda-1)(f-1)^2}{r^2} - 2w(1-f-rf') \right) \quad (10)$$

with $w = 8\mu^2(3\lambda-1)/\kappa^2$. For $\lambda = 1$ ($w = 16\mu^2/\kappa^2$), we have a solution where f and N are determined to be

$$N^2 = f = \frac{2(r^2 - 2Mr + \alpha)}{r^2 + 2\alpha + \sqrt{r^4 + 8\alpha Mr}}, \quad (11)$$

where $\alpha = 1/(2w)$ and M is an integration constant related to the mass. The metric of this black hole looks like that of Gauss-Bonnet black hole. The event horizons are given by

$$r_{\pm} = M \pm \sqrt{M^2 - \alpha}, \quad (12)$$

and the Hawking temperature is

$$T_H = \frac{f'}{4\pi} \Big|_{r=r_+} = \frac{\sqrt{M^2 - \alpha}}{\pi(r_+^2 + 2\alpha + \sqrt{r_+^4 + 8\alpha Mr_+})}. \quad (13)$$

The thermodynamics of this black hole has been studied in [28, 30, 31]. Myung [30] has calculated the ADM mass of this black hole and find that it is similar to that of 4-dimensional Einstein-Born-Infeld black hole [32]. It is well known that the Born-Infeld electrodynamics is one of the important nonlinear electromagnetic theories. As the Born-Infeld scale parameter b tends to zero, the Einstein-Maxwell theory is recovered and the Einstein-Born-Infeld black hole is reduced to Reissner-Norström black hole. Applying such kind of ADM mass proposed in [30], Wang *et al* [31] find that the integral and differential forms of the first law of thermodynamics are still valid for the deformed Hořava-Lifshitz black hole. The potentially observable properties of this black hole were considered in [33–35]. In this letter, our main purpose is to study the quasinormal modes of massless scalar field in this spacetimes and to see what there exists some new feature in the dynamical evolution of the perturbation in the black hole in Hořava-Lifshitz theory.

The Klein-Gordon equation for a massless scalar field in this spacetime is

$$\frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu) \psi = 0. \quad (14)$$

Separating $\psi = e^{-i\omega t} R(r) Y_{lm}(\theta, \phi)/r$, we can obtain the radial equation for the scalar perturbation in the deformed Hořava-Lifshitz black hole spacetime

$$\frac{d^2 R(r)}{dr_*^2} + [\omega^2 - V(r)] R(r) = 0, \quad (15)$$

where r_* is the tortoise coordinate (which is defined by $dr_* = \frac{1}{f}dr$) and the effective potential $V(r)$ reads

$$V(r) = \frac{2(r^2 - 2Mr + \alpha)}{r^2 + 2\alpha + \sqrt{r^4 + 8\alpha Mr}} \left[\frac{l(l+1)}{r^2} + \frac{1}{\alpha} \left(1 - \frac{r^3 + 2\alpha M}{r\sqrt{r^4 + 8\alpha Mr}} \right) \right]. \quad (16)$$

Obviously, as $\alpha = 0$ the effective potential $V(r)$ can be reduced to that of the Schwarzschild black hole. When α increases we find that the peak value of the potential barrier gets lower for $l = 0$ and higher for $l = 1$, which is shown in figure (1). In the Reissner-Norström black hole background, for all l the peak value of the effective potential increases with the charge q of the black hole. This means that although the formulas of the outer and inner horizons are very similar, the behaviors of the effective potential are different in these black hole background. In the Einstein-Born-Infeld black hole, the variety of the effective potential with the charge q is similar to that in Reissner-Norström black hole spacetime. However, the Born-Infeld scale parameter b decreases the peak value of $V(r)$ for all l . These results imply the quasinormal modes in the deformed Hořava-Lifshitz black hole possess some different properties from those of the black holes in the Einstein-Maxwell and Einstein-Born-Infeld gravities.




FIG. 1: Variety of the effective potential $V(r)$ with the polar coordinate r in the deformed Hořava-Lifshitz black hole (left column), Reissner-Norström black hole (middle column) and Einstein-Born-Infeld black hole with fixed scale parameter $b = 1.0$ (right column). The figures in the upper row are for $l = 0$ and in the lower row are for $l = 1$ and $M = 1$.

We are now in a position to apply the the third-order WKB approximation method approximation to

evaluate the fundamental quasinormal modes ($n = 0$) of massless scalar perturbation in the deformed Hořava-Lifshitz black hole. We expect to see what effects of Hořava-Lifshitz parameter α can be reflected in the quasinormal modes' behavior. The formula for the complex quasinormal frequencies ω in this approximation is given by [36–38]

$$\omega^2 = [V_0 + (-2V_0'')^{1/2}\Lambda] - i(n + \frac{1}{2})(-2V_0'')^{1/2}(1 + \Omega), \quad (17)$$

where

$$\begin{aligned} \Lambda &= \frac{1}{(-2V_0'')^{1/2}} \left\{ \frac{1}{8} \left(\frac{V_0^{(4)}}{V_0''} \right) \left(\frac{1}{4} + \alpha^2 \right) - \frac{1}{288} \left(\frac{V_0'''}{V_0''} \right)^2 (7 + 60\alpha^2) \right\}, \\ \Omega &= \frac{1}{(-2V_0'')^{1/2}} \left\{ \frac{5}{6912} \left(\frac{V_0'''}{V_0''} \right)^4 (77 + 188\alpha^2) \right. \\ &\quad - \frac{1}{384} \left(\frac{V_0'''^2 V_0^{(4)}}{V_0''^3} \right) (51 + 100\alpha^2) + \frac{1}{2304} \left(\frac{V_0^{(4)}}{V_0''} \right)^2 (67 + 68\alpha^2) \\ &\quad \left. + \frac{1}{288} \left(\frac{V_0'''}{V_0''} \right) \left(\frac{V_0^{(5)}}{V_0''^2} \right) (19 + 28\alpha^2) - \frac{1}{288} \left(\frac{V_0^{(6)}}{V_0''} \right) (5 + 4\alpha^2) \right\}, \end{aligned} \quad (18)$$

and

$$\alpha = n + \frac{1}{2}, \quad V_0^{(s)} = \left. \frac{d^s V}{dr_*^s} \right|_{r_* = r_*(r_p)},$$

n is overtone number and r_p is the value of polar coordinate r corresponding to the peak of the effective potential (16). Setting $M = 1$ and substituting the effective potential (16) into the formula above, we can obtain the quasinormal frequencies of scalar perturbation in the deformed Hořava-Lifshitz black hole.

α	$\omega \quad (l = 0)$	$\omega \quad (l = 1)$	$\omega \quad (l = 2)$
0.0	0.104647-0.115197i	0.291114-0.098001i	0.483211-0.096805i
0.1	0.104687-0.111855i	0.293313-0.096159i	0.486927-0.095211i
0.2	0.104300-0.108459i	0.295640-0.094238i	0.490850-0.093510i
0.3	0.103481-0.105037i	0.298101-0.092219i	0.495005-0.091681i
0.4	0.102257-0.101646i	0.300704-0.090070i	0.499417-0.089697i
0.5	0.100690-0.098374i	0.303454-0.087770i	0.504120-0.087522i

TABLE I: The fundamental ($n = 0$) quasinormal frequencies of scalar field in the in the deformed Hořava-Lifshitz black hole for $l = 0, 1, 2$.

In tables (I)-(IV), we list the fundamental quasinormal frequencies of the massless scalar perturbation field for fixed $l = 0, 1$ and 2 in the deformed Hořava-Lifshitz, the Reissner-Norström and Einstein-Born-Infeld black hole spacetimes, respectively. From the table (I) and figures (2) and (3), we find that with the increase of the parameter α the real parts decrease for $l = 0$ and increase for $l = 1$ and $l = 2$. The absolute value

q	$\omega \quad (l=0)$	$\omega \quad (l=1)$	$\omega \quad (l=2)$
0.0	0.104647-0.115197i	0.291114-0.098001i	0.483211-0.096805i
0.1	0.104856-0.115204i	0.291616-0.098051i	0.484025-0.096857i
0.2	0.105493-0.115221i	0.293145-0.098196i	0.486503-0.097011i
0.3	0.106586-0.115230i	0.295770-0.098430i	0.490758-0.097261i
0.4	0.108179-0.115195i	0.299617-0.098737i	0.496997-0.097593i
0.5	0.110339-0.115049i	0.304891-0.099084i	0.505561-0.097977i

TABLE II: The fundamental ($n = 0$) quasinormal frequencies of scalar field in the Reissner-Norström black hole for $l = 0, 1, 2$.

q	$\omega \quad (l=0)$	$\omega \quad (l=1)$	$\omega \quad (l=2)$
0.0	0.104647-0.115197i	0.291114-0.0980010i	0.483211-0.0968050i
0.1	0.104786-0.115202i	0.291449-0.0980342i	0.483753-0.0968396i
0.2	0.105209-0.115215i	0.292462-0.0981318i	0.485395-0.0969430i
0.3	0.105927-0.115228i	0.294184-0.0982911i	0.488185-0.0971125i
0.4	0.106961-0.115229i	0.296667-0.0985064i	0.492212-0.0973430i
0.5	0.108344-0.115194i	0.299995-0.0987669i	0.497609-0.0976251i

TABLE III: The fundamental ($n = 0$) quasinormal frequencies of scalar field in the Einstein-Born-Infeld black hole with $b = 1.0$ for $l = 0, 1, 2$.

of imaginary parts for all l decrease. Comparing with the quasinormal modes of the Reissner-Norström and Einstein-Born-Infeld black holes [39–42], we find from the tables (I)-(III) that the dependence of quasinormal modes on the parameter α is different from that on the charge q in the usual Reissner-Norström and Einstein-Born-Infeld black hole spacetimes. With the increase of q , the real part of the quasinormal modes increases in both Reissner-Norström and Einstein-Born-Infeld black holes. While the absolute value of imaginary parts first increase and then decrease. Moreover, from tables (I) and (IV), we also find the effect of the parameter α on quasinormal modes is different from that of the Born-Infeld parameter b in the Einstein-Born-Infeld black hole. As b increases, both the real parts and the absolute value of imaginary parts decrease for all l . Moreover, the speed of decrease of the absolute value of imaginary parts with b in the Einstein-Born-Infeld black hole is smaller than that with α in the deformed Hořava-Lifshitz black hole. The tables (I)-(IV) also tell us that the absolute value of imaginary parts is smaller than that of in the Schwarzschild, Reissner-Norström and Einstein-Born-Infeld black hole spacetimes. This means that the scalar perturbation decays more slowly in the deformed Hořava-Lifshitz black hole.

For the late-time evolution of the massless scalar perturbation, one can expect that it has the same form as that in background of a Schwarzschild black hole because that in the deformed Hořava-Lifshitz black hole spacetimes the effective potentials have the same asymptotic behaviors when the polar coordinate r approaches to infinite.

b	ω ($l=0$)	ω ($l=1$)	ω ($l=2$)
0.0	0.105493-0.115221i	0.293145-0.0981960i	0.486503-0.0970110i
0.1	0.105465-0.115221i	0.293077-0.0981895i	0.486392-0.0970044i
0.2	0.105436-0.115220i	0.293008-0.0981831i	0.486280-0.0969976i
0.3	0.105408-0.115220i	0.292940-0.0981768i	0.486169-0.0969908i
0.4	0.105379-0.115219i	0.292871-0.0981704i	0.486058-0.0969840i
0.5	0.105351-0.115218i	0.292803-0.0981640i	0.485947-0.0969772i

TABLE IV: The fundamental ($n=0$) quasinormal frequencies of scalar field in the Einstein-Born-Infeld black hole with $q=0.2$ for $l=0, 1, 2$.




FIG. 2: Variety of the real part of the fundamental quasinormal modes with α in the deformed Hořava-Lifshitz black hole spacetime. The left is for $l=0$ and the right for $l=1, 2$.

In summary we study the quasinormal modes of the massless scalar perturbation in the background of a deformed black hole in the Hořava-Lifshitz gravity. Our results show that the dependence of quasinormal modes on the parameter α is different from that on the charge q in the usual Reissner-Norström black hole spacetime. Our analysis also indicate that the effect of the parameter α on quasinormal modes is different from that of the Born-Infeld parameter b in the Einstein-Born-Infeld black hole. Moreover, we also find the absolute value of imaginary parts is smaller than that of in the Schwarzschild, Reissner-Norström and Einstein-Born-Infeld black hole spacetimes, which means that the scalar perturbation decays more slowly in the deformed Hořava-Lifshitz black hole. This would open a window to examine the Hořava-Lifshitz theory, Einstein-Maxwell theory and Einstein-Born-Infeld theory in the future. It would be of interest to generalize our study to the other Hořava-Lifshitz black hole spacetimes in which $\lambda \neq 1$, such as the black holes in refs [19, 21–24]. Work in this direction will be reported in the future.




FIG. 3: Variety of the absolute value of imaginary parts of the fundamental quasinormal modes with α for fixed $l = 1, 2$ in the deformed Hořava-Lifshitz black hole spacetime.

Acknowledgments

This work was partially supported by the National Natural Science Foundation of China under Grant No.10875041; the Scientific Research Fund of Hunan Provincial Education Department Grant No.07B043 and the construct program of key disciplines in Hunan Province. J. L. Jing's work was partially supported by the National Natural Science Foundation of China under Grant No.10675045; the FANEDD under Grant No. 200317; the National Basic Research of China under Grant No. 2010CB833004; and the Hunan Provincial Natural Science Foundation of China under Grant No.08JJ3010.

-
- [1] P. Horava, Phys. Rev. D **79** (2009) 084008.
 - [2] P. Horava, JHEP **0903** (2009) 020.
 - [3] P. Hořava, Phys.Rev.Lett.**102** (2009) 161301, arXiv:0902.3657.
 - [4] A. Volovich and C. Wen, JHEP **0905** (2009) 087, arXiv: 0903.2455.
 - [5] J. Kluson, JHEP **0907** (2009) 079, arXiv:0904.1343.
 - [6] H. Nikolic, arXiv:0904.3412.
 - [7] H. Nastase, arXiv:0904.3604.
 - [8] K. I. Izawa, arXiv:0904.3593.
 - [9] G. E. Volovik, JETP Lett.**89** (2009) 525, arXiv:0904.4113.
 - [10] B. Chen and Q. G. Huang, Phys. Lett.B **683** (2010) 108, arXiv:0904.4565.
 - [11] R. G. Cai, B. Hu and H. B. Zhang, Phys. Rev. D**80** (2009) 041501, arXiv:0905.0255.
 - [12] G. Calcagni, JHEP **0909** (2009) 112, arXiv:0904.0829.
 - [13] T. Takahashi and J. Soda, Phys. Rev. Lett.**102** (2009) 231301, arXiv:0904.0554.
 - [14] S. Mukohyama, JCAP **0906** (2009) 001, arXiv:0904.2190.
 - [15] R. Brandenberger, arXiv:0904.2835.

- [16] Y. S. Piao, arXiv:0904.4117.
- [17] X. Gao, arXiv:0904.4187.
- [18] E. Kiritsis and G. Kofinas, Nucl.Phys.B **821** (2009) 467, arXiv:0904.1334.
- [19] E. O. Colgain and H. Yavartanoo, JHEP **0908** (2009) 021, arXiv:0904.4357.
- [20] A. Kehagias and K. Sfetsos, Phys. Lett. B **678** (2009) 123, arXiv:0905.0477.
- [21] H. Lu, J. Mei and C. N. Pope, Phys. Rev. Lett. **103** (2009) 091301, arXiv:0904.1595.
- [22] R. G. Cai, L. M. Cao and N. Ohta, Phys. Rev. D **80** (2009) 024003, arXiv:0904.3670.
- [23] R. G. Cai, Y. Liu and Y. W. Sun, JHEP **0906** (2009) 010, arXiv:0904.4104.
- [24] A. Ghodsi, arXiv:0905.0836.
- [25] Y. S. Myung and Y. W. Kim, arXiv:0905.0179.
- [26] T. Nishioka, Class.Quant.Grav. **26** (2009) 242001, arXiv:0905.0473.
- [27] R. G. Cai, L. M. Cao and N. Ohta, Phys. Lett.B **679** (2009) 504, arXiv:0905.0751.
- [28] Y. S. Myung, Phys. Lett. B **678** (2009) 127, arXiv:0905.0957.
- [29] R. L. Arnowitt, S. Deser and C.W. Misner, *The dynamics of general relativity*, “Gravitation: an introduction to current research”, Louis Witten ed. (Wiley 1962), chapter 7, pp 227-265, arXiv:gr-qc/0405109.
- [30] Y. S. Myung, Phys. Lett. B **685** (2010) 318, arXiv:0912.3305.
- [31] M. Wang, J. Jing, C. Ding, S. Chen, arXiv:0912.4832.
- [32] N. Breton, Phys. Rev. D **67** (2003) 124004.
- [33] T. Harko, Z. Kovács, F. S. N. Lobo, Phys. Rev. D **80** (2009) 044021;
T. Harko, Z. Kovács, F. S. N. Lobo, arXiv:0908.2874.
- [34] R. A. Konoplya, Phys. Lett.B **679** (2009) 499.
- [35] S. B. Chen and J. L. Jing, Phys.Rev.D **80** (2009) 024036.
- [36] B. F. Schutz and C. M. Will, Astrophys. J. Lett. Ed. **291**, L33 (1985).
- [37] S. Iyer and C. M. Will, Phys. Rev. D **35**, 3621 (1987).
- [38] S. Iyer, Phys. Rev. D **35**, 3632 (1987).
- [39] R. A. Konoplya, Phys.Lett. B **550**, 117 (2002).
- [40] R. A. Konoplya, Phys. Rev. D **66**, 084007 (2002)
- [41] H. Onozawa, T. Mishima, T. Okamura and H. Ishihara, Phys. Rev. D **53**, 7033 (1996).
- [42] S. Fernando, C. Holbrook, Int. J. Theor.Phys. **45** (2006) 1630; S. Fernando, Gen. Rel. Grav. **37** (2005) 585.